

On discrepancy between ATIC and Fermi data

Dmitry Malyshev*

CCPP, 4 Washington Place, Meyer Hall of Physics, NYU, New York, NY 10003

(Dated: July 4, 2009)

Either ATIC or Fermi-LAT data can be fitted together with the PAMELA data by three components: primary background $\sim E^{-3.3}$, secondary background $\sim E^{-3.6}$, and an additional source of electrons $\sim E^{-\gamma_a} \text{Exp}(-E/E_{\text{cut}})$. We find that the best fits for ATIC + PAMELA and for Fermi + PAMELA are approximately the same, $\gamma_a \approx 2$ and $E_{\text{cut}} \sim 500$ GeV. However, the ATIC data have a narrow bump between 300 GeV and 600 GeV which contradicts the smooth Fermi spectrum. An interpretation of the ATIC bump as well as the featureless Fermi spectrum in terms of dark matter models and pulsars is discussed.

*Electronic address: dm137@nyu.edu; On leave of absence from ITEP, Moscow, Russia, B. Cheremushkinskaya

The question of interpretation of Fermi-LAT [1], HESS [2, 3], and ATIC data [4] can be split into two parts: general properties of the flux and the presence of features. If one takes into account PAMELA data [5], then both Fermi and ATIC require the existence of an additional flux of electrons and positrons complementary to the standard primary and secondary backgrounds.

We will consider the following form for the additional flux

$$F_a \sim E^{-\gamma_a} e^{-\frac{E}{E_{cut}}}. \quad (1)$$

The general properties of the flux will be parameterized by the index γ_a and the exponential cutoff E_{cut} . Significant deviations from this form will be considered as “features”.

We will assume the primary background $\sim E^{-\gamma_p}$, the index $\gamma_p \approx 3.3$ can be estimated from the electron injection index $2 - 2.5$ due to shock acceleration in the supernovae explosions [6, 7] (a similar estimation for the shock acceleration in gamma ray bursts can be found in [8, 9]). The shift of the index from $2 - 2.5$ to 3.3 is due to cooling during propagation [10]. Secondary background of electrons and positrons is produced by collisions of high energy protons and nuclei $\sim E^{-2.7}$ [11, 12] with the dust. We will assume the secondary background $\sim E^{-\gamma_s}$, $\gamma_s \approx 3.6$ where the difference in the indices is again due to cooling. Since the normalization and the indices of the backgrounds are known only approximately, we will treat them as independent parameters in the fits.

In order to find the properties of the additional flux, we will use the ATIC data, the Fermi data, and the PAMELA data above 10 GeV (the points below 10 GeV are assumed to suffer from solar modulation and we will discard them for the purposes of current analysis). The results of the fits to ATIC + PAMELA and to Fermi + PAMELA are presented in Figs. 1 and 2 and in Table I. Both ATIC + PAMELA and Fermi + PAMELA are fitted best by an additional flux with $\gamma_a \approx 2$ and $E_{cut} \sim 500$ GeV, i.e., these experiments are consistent with each other from the point of view of general properties of the flux parameterized by Eq. (1). It should be noted that without PAMELA data, the ATIC bump is better fitted with a harder additional flux, $\gamma_a \approx 1.7$, while the Fermi data are consistent with the primary background $\gamma_p \approx 3.0 - 3.1$ [1, 13].

The second question is the presence of features. The Fermi data are well approximated by the additional source in Eq. (1) while the ATIC data have a deviation from (1) between 300 GeV and 600 GeV. Thus, although the parameters of the additional flux in Eq. (1) necessary to fit ATIC and Fermi data are similar, the ATIC data have a bump while the Fermi data don't have any features, i.e., there is a disagreement between the two data sets at energies 300 - 600 GeV. As one can see in Table I, the best fit for Fermi + PAMELA has a better reduced chi-squared, $\chi_r^2 = 0.4$, than the best fit for ATIC + PAMELA, $\chi_r^2 = 1.5$.

In the following we will summarize the properties of dark matter (DM) models and pulsars necessary to reproduce both the flux with features and the flux without features. Due to energy losses, the source of high energy electrons should be close to Earth, within approximately 1 - 3




FIG. 1: The results of fitting the primary and secondary backgrounds together with an additional source of electrons and positrons to Fermi + PAMELA and to ATIC + PAMELA. As everywhere else in the paper, for PAMELA, only $E > 10$ GeV points are used. The best fit values and the parameter ranges are given in Table I. The star (cross) represents the best fit to Fermi + PAMELA (ATIC + PAMELA) with the reduced chi-squared $\chi_{\text{r best}}^2 = 0.4$ ($\chi_{\text{r best}}^2 = 1.5$). The dashed (solid) contours correspond to $\chi_{\text{r best}}^2 + 1$ and $\chi_{\text{r best}}^2 + 2$ for Fermi (ATIC) + PAMELA. The concordance model has $\chi_{\text{r}}^2 = 0.9$ (2.0) for Fermi (ATIC) + PAMELA.

kpc. For the purposes of flux calculation, the DM distribution can be viewed as homogeneous and constant [14] (unless there is a significant contribution from a local DM substructure such as a clump [14, 15, 16]). The flux from a homogeneous source is [10]

$$F_{\text{DM}} = \frac{c}{4\pi} \frac{1}{b(E)} \int_E^{M_{\text{DM}}} Q_{\text{DM}}(\tilde{E}) d\tilde{E}, \quad (2)$$

where $\dot{E} \equiv -b(E)$ is the energy losses. At $E > 10$ GeV the energy losses are due to Inverse Compton Scattering and synchrotron radiation in the galactic magnetic field, thus $b(E) = b_0 E^2$ [10]. Q_{DM} is the source function for e^+e^- produced by annihilating or decaying DM, $Q_{\text{DM}} = \frac{dN}{dEdVdt}$.

The cutoff energy E_{cut} is the energy where the integral on the right hand side of Eq. (2) is saturated. For energies $E \ll E_{\text{cut}}$ the integral is insensitive to the variations of the lower limit and $F_{\text{DM}} \sim 1/b(E)$. Thus an index $\gamma_a \approx 2$ is a universal prediction of DM. Local clumps of DM result in a harder spectrum $\gamma_a < 2$ [14, 15]. The dependence on the host halo profile leads to a softer spectrum $\gamma_a > 2$ at low energies [14]. In models with small diffusion height, there may be some hardening $\gamma_a < 2$ at low energies due to leakage of electrons from the Galaxy (see, e.g., model M2 in Figure 3 of [17]).

For a given $E_{\text{cut}} \sim 500$ GeV, the DM mass and the shape of the spectrum depend on the DM

Data	γ_a	E_{cut} (GeV)	$E_0^3 F(E_0)$ (GeV ² m ⁻² s ⁻¹ sr ⁻¹)	χ_r^2	γ_p	γ_s
ATIC + PAMELA	1.95 ± 0.15	480 ± 200	50 ± 8	$1.5 - 2.5$	3.27	3.63
Fermi + PAMELA	2.15 ± 0.10	500 ± 150	44 ± 6	$0.4 - 1.4$	3.31	3.63
Concordance model	2.05	450	47	1.3	3.29	3.63

TABLE I: Numerical values for the fits presented in Fig. 1. The normalization is given for $E_0 = 100$ GeV. The ranges of parameters correspond to the ranges of reduced chi-squared χ_r^2 in the fifth column. The error bars for Fermi are computed as square root of systematic plus statistical errors squared. γ_p and γ_s are the indices of the primary and secondary backgrounds respectively. The χ_r^2 for the concordance model is computed using Fermi, ATIC, and PAMELA (>10 GeV) points. The best fits are found by varying 7 parameters: 2 indices and 2 normalization constants for primary and secondary backgrounds together with the index, the normalization and the cutoff of the additional flux in Eq. (1).

model. In general, DM annihilation is followed by a sequence of decays leading to electrons and positrons together with other stable particles in the end. Models with many steps in the decay process have smooth e^+e^- spectrum and a large DM mass $M_{\text{DM}} \gg E_{\text{cut}}$ [14, 18]. These models are favored [19, 20] by featureless Fermi spectrum. Models with decay channels through W , Z gauge bosons and quarks have stricter constraints due to absence of significant deviations from the expected backgrounds for anti-protons [21] and diffuse gamma rays from the Galactic center [22, 23]. The flux of gamma rays from DM clumps assuming $b\bar{b}$ annihilation channel was also estimated [24]. DM models with small DM mass $M_{\text{DM}} \lesssim 500$ GeV (e.g., [25, 26]) seem to be in tension with Fermi data due to absence of significant step-like features below or around 500 GeV.

DM models with few decay steps have a sharper cutoff and $M_{\text{DM}} \gtrsim E_{\text{cut}}$ [14, 18, 27]. These models give reasonable fits to the Fermi data for $M_{\text{DM}} \lesssim 1$ TeV [13], however, due to a sharp cutoff near M_{DM} they, generally, fit better the ATIC data [18]. These models may have additional constrained due to final state radiation from the Galactic center and the Galactic ridge [18, 27].

Local clumps may produce additional features at high energies. The presence of a large local clump is disfavored by Fermi but consistent with ATIC [14]. Furthermore, in models with many decay steps, a significant contribution from a local clump may be necessary to fit the ATIC bump.

Thus, the Fermi data favor dark matter with many decay steps and a large DM mass while ATIC requires either a DM model with few decay steps or a significant local substructure in DM density distribution.

Let us now turn to pulsars. In the calculation of electron and positron fluxes, pulsars can be considered as point-like instantaneous sources [28], $Q \sim \delta(x - x_0)\delta(t - t_0)$. The main reason is that the typical propagation time ($\gtrsim 100$ kyr) is much larger than the characteristic time scale




FIG. 2: The concordance model corresponding to the triangle in Figure 1 with the parameters given in Table I. The upper (lower) edge of the band is the best fit for ATIC (Fermi) + PAMELA points with energies $E > 10$ GeV. HESS 2008 and HESS 2009 points fit the concordance model, if multiplied by a factor ~ 0.9 which may be related to contamination by diffuse gamma rays.

when a pulsar loses most of its rotational energy and the electrons and positrons are released to the interstellar medium (ISM).

For every pulsar, we will consider two energy scales [13, 28]: the cutoff in the injection spectrum of electrons and positrons from the pulsar into the ISM, $E_{\text{inj.cut}}$, which can be between few hundred GeVs and tens of TeVs (see, e.g., [29] and references in [28]), and the cooling break $E_{\text{br}} = \frac{1}{b_0 t}$, obtained by integrating the energy losses $\dot{E} = -b_0 E^2$ during the propagation time t , where t is approximately the age of the pulsar. At $E \approx 100$ GeV the energy losses can be

estimated as $b_0 \approx 1.6 \text{ GeV}^{-1} \text{ s}^{-1} \approx 5 \text{ TeV}^{-1} \text{ Myr}^{-1}$ [30]. At energies $E \gtrsim 100 \text{ GeV}$ the coefficient b_0 slightly decreases with the energy [30], since the Thompson approximation to inverse Compton scattering between electrons and the starlight becomes inapplicable.

For pulsars with age $t \lesssim 10 \text{ Myr}$, the cooling break is $E_{\text{br}} \gtrsim 20 \text{ GeV}$. In the ATNF catalog there are several hundred pulsars within 3 kpc from Earth and an age $t < 10 \text{ Myr}$ [31]. Below approximately 300 GeV, the corresponding flux is well approximated by the flux from a continuous distribution of pulsars in the Galactic plane [28]. An index $\gamma_a \approx 2$ requires an index in the electron injection spectrum from pulsars $\gamma_{\text{inj}} \lesssim 2$ [28, 32, 33, 34, 35, 36, 37].

At energies $E \gtrsim 300 \text{ GeV}$, the flux receives contributions only from young pulsars $t \lesssim 1 \text{ Myr}$ within a smaller distance $d \lesssim 1 \text{ kpc}$ (since the propagation time is smaller). One may expect only of order ten such pulsars [31]. There is also a lower bound on the age of pulsars, a few tens of kyr, due to the fact that the electrons may still be trapped by the Pulsar Wind Nebulae.

Depending on the relative value of $E_{\text{inj.cut}}$ and E_{cut} , there are two possibilities for the e^+e^- flux from pulsars:

- $E_{\text{inj.cut}} \sim E_{\text{cut}}$, then the observed spectrum is naturally flat if we assume that the injection spectrum from pulsars is flat. This possibility is favored by the Fermi data [13].
- $E_{\text{inj.cut}} \gg E_{\text{cut}}$, then the cutoff in the observed spectrum is due to the cooling break which is much sharper than an exponential cutoff. One should also expect a series of steps due to consecutive cooling break cutoffs from different pulsars [28]. This possibility is consistent with ATIC but may be in tension with Fermi.

To summarize, both Fermi and ATIC require an additional source of electrons and positrons with an index $\gamma_a \approx 2$ at low energies and a cutoff $E_{\text{cut}} \sim 500 \text{ GeV}$. However the presence of a bump at high energies in ATIC data contradicts the smooth spectrum of Fermi and HESS. The sources that produce featureless spectrum include the DM models with $M_{\text{DM}} > 1 \text{ TeV}$ and several steps in DM annihilation process as well as pulsars with an injection cutoff $E_{\text{inj.cut}} \sim E_{\text{cut}}$. The sources that can produce ATIC bump include DM models with $M_{\text{DM}} \sim E_{\text{cut}}$ and few decay steps or DM models with $M_{\text{DM}} \gg E_{\text{cut}}$ and a significant contribution from a local clump. Pulsars with $E_{\text{inj.cut}} \gg E_{\text{cut}}$ may give a sharp cutoff in the observed e^+e^- flux and, possibly, step-like features due to a series of cooling break cutoffs from the youngest nearby pulsars. This possibility is in tension with the Fermi data but consistent with the ATIC.

Acknowledgments.

The author is thankful to Mirko Boezio, Ilias Cholis, Joseph Gelfand, Lisa Goodenough, Michael Kuhlen, Neal Weiner, and Weiqun Zhang for valuable discussions. This work is supported in part by the Russian Foundation of Basic Research under grant RFBR 09-02-00253 and by the NSF grants PHY-0245068 and PHY-0758032.

-
- [1] **The Fermi LAT** Collaboration, A. A. Abdo *et. al.*, *Measurement of the Cosmic Ray e^+ plus e^- spectrum from 20 GeV to 1 TeV with the Fermi Large Area Telescope*, [arXiv:0905.0025](#).
 - [2] **H.E.S.S.** Collaboration, F. Aharonian *et. al.*, *The energy spectrum of cosmic-ray electrons at TeV energies*, *Phys. Rev. Lett.* **101** (2008) 261104, [[arXiv:0811.3894](#)].
 - [3] **H.E.S.S.** Collaboration, F. Aharonian *et. al.*, *Probing the ATIC peak in the cosmic-ray electron spectrum with H.E.S.S.*, [arXiv:0905.0105](#).
 - [4] J. Chang *et. al.*, *An excess of cosmic ray electrons at energies of 300.800 GeV*, *Nature* **456** (2008) 362–365.
 - [5] O. Adriani *et. al.*, *Observation of an anomalous positron abundance in the cosmic radiation*, [arXiv:0810.4995](#).
 - [6] D. C. Ellison and S. P. Reynolds, *Electron acceleration in a nonlinear shock model with applications to supernova remnants*, *Astrophys. J.* **382** (1991) 242–254.
 - [7] S. P. Reynolds and D. C. Ellison, *Electron acceleration in Tycho’s and Kepler’s supernova remnants - Spectral evidence of Fermi shock acceleration*, *Astrophys. J.* **399** (1992) L75–L78.
 - [8] A. Achterberg, Y. A. Gallant, J. G. Kirk, and A. W. Guthmann, *Particle acceleration by ultra-relativistic shocks: Theory and simulations*, *Mon. Not. Roy. Astron. Soc.* **328** (2001) 393, [[astro-ph/0107530](#)].
 - [9] T. Piran, *The physics of gamma-ray bursts*, *Rev. Mod. Phys.* **76** (2004) 1143–1210, [[astro-ph/0405503](#)].
 - [10] M. S. Longair, *High-energy astrophysics*. Cambridge University Press, 1992.
 - [11] A. W. Strong, I. V. Moskalenko, and V. S. Ptuskin, *Cosmic-ray propagation and interactions in the Galaxy*, *Ann. Rev. Nucl. Part. Sci.* **57** (2007) 285–327, [[astro-ph/0701517](#)].
 - [12] **Particle Data Group** Collaboration, C. Amsler *et. al.*, *Review of particle physics*, *Phys. Lett.* **B667** (2008) 1.
 - [13] **FERMI-LAT** Collaboration, D. Grasso *et. al.*, *On possible interpretations of the high energy electron-positron spectrum measured by the Fermi Large Area Telescope*, [arXiv:0905.0636](#).
 - [14] M. Kuhlen and D. Malyshev, *ATIC, PAMELA, HESS, Fermi and nearby Dark Matter subhalos*, [arXiv:0904.3378](#).
 - [15] D. Hooper, A. Stebbins, and K. M. Zurek, *The PAMELA and ATIC Excesses From a Nearby Clump of Neutralino Dark Matter*, [arXiv:0812.3202](#).
 - [16] P. Brun, T. Delahaye, J. Diemand, S. Profumo, and P. Salati, *The cosmic ray lepton puzzle in the light of cosmological N-body simulations*, [arXiv:0904.0812](#).
 - [17] C.-R. Chen, K. Hamaguchi, M. M. Nojiri, F. Takahashi, and S. Torii, *Dark Matter Model Selection and the ATIC/PPB-BETS anomaly*, [arXiv:0812.4200](#).

- [18] J. Mardon, Y. Nomura, D. Stolarski, and J. Thaler, *Dark Matter Signals from Cascade Annihilations*, [arXiv:0901.2926](#).
- [19] P. Meade, M. Papucci, A. Strumia, and T. Volansky, *Dark Matter Interpretations of the Electron/Positron Excesses after FERMI*, [arXiv:0905.0480](#).
- [20] L. Bergstrom, J. Edsjo, and G. Zaharijas, *Dark matter interpretation of recent electron and positron data*, [arXiv:0905.0333](#).
- [21] O. Adriani *et. al.*, *A new measurement of the antiproton-to-proton flux ratio up to 100 GeV in the cosmic radiation*, [arXiv:0810.4994](#).
- [22] **H.E.S.S.** Collaboration, F. Aharonian *et. al.*, *Discovery of Very-High-Energy Gamma-Rays from the Galactic Centre Ridge*, *Nature* **439** (2006) 695–698, [[astro-ph/0603021](#)].
- [23] **H.E.S.S.** Collaboration, F. Aharonian *et. al.*, *HESS observations of the galactic center region and their possible dark matter interpretation*, *Phys. Rev. Lett.* **97** (2006) 221102, [[astro-ph/0610509](#)].
- [24] M. Kuhlen, J. Diemand, and P. Madau, *The Dark Matter Annihilation Signal from Galactic Substructure: Predictions for GLAST*, [arXiv:0805.4416](#).
- [25] L. Bergstrom, T. Bringmann, and J. Edsjo, *New Positron Spectral Features from Supersymmetric Dark Matter - a Way to Explain the PAMELA Data?*, *Phys. Rev.* **D77** (2008) 103520, [[arXiv:0808.3725](#)].
- [26] P. Grajek, G. Kane, D. Phalen, A. Pierce, and S. Watson, *Is the PAMELA Positron Excess Winos?*, [arXiv:0812.4555](#).
- [27] P. Meade, M. Papucci, and T. Volansky, *Dark Matter Sees The Light*, [arXiv:0901.2925](#).
- [28] D. Malyshev, I. Cholis, and J. Gelfand, *Pulsars versus Dark Matter Interpretation of ATIC/PAMELA*, [arXiv:0903.1310](#).
- [29] J. Arons, *Pulsars as Gamma-Rays Sources: Nebular Shocks and Magnetospheric Gaps*, *Space Sci. Rev.* **75** (1996) 235–255.
- [30] T. Kobayashi, Y. Komori, K. Yoshida, and J. Nishimura, *The most likely sources of high energy cosmic-ray electrons in supernova remnants*, *Astrophys. J.* **601** (2004) 340–351, [[astro-ph/0308470](#)].
- [31] R. N. Manchester, G. B. Hobbs, A. Teoh, and M. Hobbs, *The ATNF Pulsar Catalogue*, [astro-ph/0412641](#). <http://www.atnf.csiro.au/research/pulsar/psrcat>.
- [32] D. Hooper, P. Blasi, and P. D. Serpico, *Pulsars as the Sources of High Energy Cosmic Ray Positrons*, *JCAP* **0901** (2009) 025, [[arXiv:0810.1527](#)].
- [33] H. Yuksel, M. D. Kistler, and T. Stanev, *TeV Gamma Rays from Geminga and the Origin of the GeV Positron Excess*, [arXiv:0810.2784](#).
- [34] S. Profumo, *Dissecting Pamela (and ATIC) with Occam's Razor: existing, well-known Pulsars naturally account for the 'anomalous' Cosmic-Ray Electron and Positron Data*, [arXiv:0812.4457](#).
- [35] K. Ioka, *A Gamma-Ray Burst/Pulsar for Cosmic-Ray Positrons with a Spectral Cutoff and Line*,

arXiv:0812.4851.

- [36] N. J. Shaviv, E. Nakar, and T. Piran, *Natural explanation for the anomalous positron to electron ratio with supernova remnants as the sole cosmic ray source*, arXiv:0902.0376.
- [37] N. Kawanaka, K. Ioka, and M. M. Nojiri, *Cosmic-Ray Electron Excess from Pulsars is Spiky or Smooth?: Continuous and Multiple Electron/Positron injections*, arXiv:0903.3782.